

# 3D Geometry for Panorama

Jianxiong Xiao  
Massachusetts Institute of Technology


## Abstract

*This document briefly explains the geometry of full-view panorama image we used in [7].*


Here, we describe how to reconstruct the rough 3D model for each single spherical image. As in [2, 4, 1], we assume that a scene is composed of a number of objects standing on a ground plane (dubbed standing objects), with the objects represented as piecewise-connected planes oriented orthogonally to the ground plane. To obtain the ground line for each spherical image, we ask people to label the ground plane polygon using LabelMe [4].

As shown in Figure 1(a), let's denote the resolution of the spherical image to be  $W \times H$  pixels and represent the panorama in Equirectangular projection [6, 5]. Because the spherical images covers 360 degree field of view horizontally and 180 degree field of view vertically, we can know that  $W = 2H$ , and the focal length is  $W/2\pi$  pixels. For spherical projection images, because of the full coverage of field of view, people normally put the camera in a position that the top of the projection sphere is pointing to the sky. Therefore, we can safely assume that the horizontal vanishing line of the ground plane is at 0 height of the image coordinate  $[-\frac{W}{2}, \frac{W}{2}] \times [-\frac{H}{2}, \frac{H}{2}]$ . As in [1, 3, 4], we further assume that the camera is at the height of  $c^h = 1.7$  meter away from the ground plane.

With these conditions, the geometry transformation for 3D reconstruction becomes very straightforward and elegant. For a 3D point  $\mathbf{P} = (X, Y, Z)$  on the ground plane with image coordinate  $(x, y)$ , we have  $Y = 0$  since it is on the ground. And  $\theta_x = 2\pi x/W$ ,  $\theta_y = \pi y/H$ . From Figure 1(b), we can see  $r = c^h |\cot \theta_y|$ . Finally, from Figure 1(c), we can see that  $X = r \cos(-\theta_x + \frac{\pi}{2})$  and  $Z = r \sin(-\theta_x + \frac{\pi}{2})$ . For a 3D point  $\mathbf{P}' = (X', Y', Z')$  not on the ground plane with image coordinate  $(x, y')$ , which is assumed to be on a dubbed standing billboard plane, we first find out a point  $\mathbf{P} = (X, Y, Z)$  on the ground plan with same  $x$  image coordinate, then we have  $X' = X$ ,  $Z' = Z$ , and  $r' = r = c^h |\cot \theta_y|$ . The only difference is  $Y' \neq Y = 0$ . From Figure 1(b), we can see that  $Y' = c^h + r \tan \theta_{y'}$ .



(a) Spherical Image Plane.



(b) Slice of 3D to calculate  $r$  and (c)  $X$  and  $Z$  are calculated from  $r$  and  $Y$ .

Figure 1. Geometry derivation for 3D reconstruction of spherical image using ground line.

## References

- [1] D. Hoiem, A. Efros, and M. Hebert. Putting objects in perspective. In *CVPR*, 2006. 1
- [2] D. Hoiem, A. A. Efros, and M. Hebert. Geometric context from a single image. In *ICCV*, 2005. 1
- [3] D. Hoiem, A. A. Efros, and M. Hebert. Closing the loop on scene interpretation. In *CVPR*, 2008. 1
- [4] B. Russell and A. Torralba. Building a database of 3d scenes from user annotations. In *CVPR*, 2009. 1
- [5] J. P. Snyder. *Flattening the Earth: Two Thousand Years of Map Projections*. 1993. 1
- [6] Wikipedia. Equirectangular projection, 2004. [http://en.wikipedia.org/wiki/Equirectangular\\_projection](http://en.wikipedia.org/wiki/Equirectangular_projection). 1
- [7] J. Xiao, K. Ehinger, A. Oliva, and A. Torralba. Recognizing scene viewpoint using panoramic place representation. In *CVPR*, 2012. 1